

ORIGINAL RESEARCH

Comparative study of USG and CT in the evaluation of suspicious Ovarian Masses

¹Dr. Priyanka, ²Dr. Shaurya Thakran

^{1,2}Assistant Professor, Department of Radiology, Gouri Devi Institute of Medical Sciences, Durgapur, West Bengal, India

Corresponding Author

Dr. Shaurya Thakran

Assistant Professor, Department of Radiology, Gouri Devi Institute of Medical Sciences, Durgapur, West Bengal, India

Received: 11 July, 2020

Accepted: 16 August, 2020

ABSTRACT

Aim: comparison of Ultrasonography (USG) and Computed Tomography (CT) in the evaluation of suspicious Ovarian Masses. **Materials and Methods:** This prospective observational study was carried out in the department of radiology. All patients underwent abdominal Ultrasonography and CT scan with determination of the ovarian mass characteristics. **Results:** There is total 30 cases of Pre-menopausal stage and 20 cases of Post-menopausal stage having ovarian cyst. Out of 30 cases of Pre-menopausal conditions have 5 number of malignant and 25 number of benign type of ovarian masses. In the Postmenopausal group there are 15 cases of malignant and 5 cases of benign ovarian mass was observed. Table 3 shows CT and USG comparison for the diagnosis of ovarian masses. Overall, CT was found to have 98% sensitivity, 91% specificity, and an accuracy of 96% in the differentiation of benign and malignant ovarian masses, while PPV and NPV were 97% and 93%, respectively. The sensitivity of USG was 88%, specificity was 86% and PPV and NPV were 87% and 83% respectively. **Conclusion:** In this present study showed significant differences in the two methods i.e USG and CT. CT is showing more advantages regarding tumor localization, characterization. Hence CT can be advised if the unusual abnormalities were observed in routine USG scan in the diagnosis of ovarian masses.

Keywords: USG, CT Ovarian Masses

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

INTRODUCTION

Adnexal lesions especially ovarian masses are a common presentation amongst women of all age groups and all social strata. Pertaining to their wide spectrum of diagnostic variation, they often perplex both the physician and the radiologist. While the docile benign ovarian lesions may be treated conservatively, the aggressive neoplastic lesions often require radical surgical and associated oncological treatment. Ovarian cancer is a silent killer as it is conspicuous by its late diagnosis and low 5 years survival rate of 45%. It is second only to cervical cancer in gynecological malignancies in India and has a worldwide prevalence.^{1,2} Ovarian cyst is often asymptomatic and it is a fluid-filled sac inside the ovary. Sometimes it leads to lower abdominal or back pain, pelvic inflammatory disease. But most of the ovarian cysts are not harmful.³ Ovarian cyst can be follicular, corpus luteum, dermoid and cystadenomas type.⁴ The diagnosis of ovarian cyst can be performed by the use of ultrasound and other laboratory investigations.⁵⁻⁸ Sometimes if required patients can take medications like ibuprofen or paracetamol. Surgical procedures can be taken in case of larger

cysts.^{9,10} Most of the reproductive age female can develop smaller cyst every month. Larger cyst can cause problems before menopause in 8% of women.¹¹ 16% of female with ovarian cyst has risk of ovarian cancer. Therefore, radiological evaluation of ovarian masses is pivotal in making early diagnosis and lesion characterization, distinguishing between benign and malignant masses thereby determining the therapeutic approach. Various diagnostic modalities such as USG, CT and now MRI have come to the rescue of the diagnostician for solving these dilemmas.¹² USG is typically the first study to be requested in patients with clinical findings that may suggest ovarian mass. The advantages of a USG are its wide availability, low cost and accuracy for morphological characterization. However, a considerable percentage of the ovarian masses may be considered as indeterminate on USG.¹³ It is for such lesions that cross-sectional imaging techniques are pivotal. MRI can provide precise anatomical localization and meticulous lesion characterization; thereby significantly narrowing down the differential diagnosis. However, in a country like India, especially in the remote locations, availability and cost

effectiveness are major issues that are preventing MRI to be the second line modality after USG for evaluating ovarian masses. CT on the other hand has wide availability, relative cost effectiveness, rapidity and provides a larger field of view allowing comprehensive evaluation of the abdomen.^{14,15}

MATERIAL AND METHODS

This prospective observational study was carried out in the department of radiology, after taking the approval of the protocol review committee and institutional ethics committee. After taking informed consent detailed history was taken from the patient or the relatives. Total 50 women were included as subjects in this prospective study. The aim and the objective of the study are conveyed to all patients. All patients underwent abdominal Ultrasonography and CT scan with determination of the ovarian mass characteristics. Patients with conservatively manageable ovarian masses were excluded from this study. Patients of age 0 to 18 years, midline uterine mass lesions on USG, clinically and sonographically proven cases of ectopic pregnancy, sonographically validated benign cystic ovarian lesions such as functional cysts in patients of reproductive age group were excluded from the study. Complete history of

allergy was taken before doing CT scan and if there was history of allergy then nonionic contrast was used.

RESULTS

We evaluated 50 patients with Mean age of 41.22 years. Most patients belong to 40-50 year age group and followed by 30-40 years. Table 1. The table 2 shows the Benign and Malignant Masses on Histopathology in Pre and Post-menopausal patients. There are total 30 cases of Pre-menopausal stage and 20 cases of Post-menopausal stage having ovarian cyst. Out of 30 cases of Pre-menopausal conditions have 5 number of malignant and 25 number of benign type of ovarian masses. In the Postmenopausal group there are 15 cases of malignant and 5 cases of benign ovarian mass was observed. Table 3 shows CT and USG comparison for the diagnosis of ovarian masses. Overall, CT was found to have 98% sensitivity, 91% specificity, and an accuracy of 96% in the differentiation of benign and malignant ovarian masses, while PPV and NPV were 97% and 93%, respectively. The sensitivity of USG was 88%, specificity was 86% and PPV and NPV were 87% and 83% respectively.

Table 1. Age distribution of patients

Age Group (in years)	Number=50	Percentage (%)
Below 20	3	6
20-30	8	16
30-40	10	20
40-50	19	38
50-60	9	18
Above 60	1	2

Table-2: The characteristics of different ovarian masses

Category	Pre-menopausal	Post-menopausal
Malignant	5	15
Benign	25	5
Total	30	20

Table-3: The comparison between USG and CT in diagnosis of ovarian masses

Category	CT Study (No. of Cases)		USG Study (No. of Cases)	
	Benign	Malignant	Benign	Malignant
Sensitivity	98%	87%	88%	77%
Specificity	91%	86%	86%	75%
Positive Predictive Value	97%	89%	87%	80%
Negative Predictive value	93%	90%	83%	75%

DISCUSSION

In day-to-day practice, we come across many cases of ovarian masses. Some of these turn out to be benign, some borderline, and some malignant. When an ovarian mass is detected, there are two major issues: to determine whether it is benign or malignant and then if it is malignant, to look for the extent of disease.^{16,17} If the nature of the mass is adequately determined on the image, then it saves the patient

unnecessary surgery and expense. Similarly, if staging is accurately done on imaging, again it becomes cost-effective and it helps in further planning.¹⁷ However, we understand that surgery has a role in definite diagnosis and the further characterization of masses. Sometimes USG underestimates staging and pelvic examination by a gynecologist and serum CA-125 are of limited value in the diagnosis of pelvic masses and their sensitivity is often below 50%.¹⁷ The sensitivity

of morphologic analysis with ultrasound in predicting malignancy in ovarian tumors has been shown to be 85%–97%, whereas its specificity ranges from 56%–95%.¹⁸⁻²¹

The above data is showing more sensitive for the detection of abnormal ovarian mass in the present population. Ovarian tumours present a greatest clinical challenge of all gynecological cancers and ovarian. Carcinoma is the second most common gynaecological carcinoma in incidence. As most of them present in a late stage, clinical diagnosis alone is difficult and as benign ovarian tumours greatly outnumber malignant ones, determination of a degree of suspicion for malignancy is critical and is based largely on imaging modalities. The determination of a degree of suspicion for malignancy in an ovarian mass is the most significant step in its management as the decision to perform radical surgery or conservative surgery depends on accurate preoperative diagnosis.¹⁹ Clinical evaluation with regards to site (unilateral or bilateral), fixity, consistency, presence of nodules in Douglas pouch and presence of ascites increase the suspicion of malignancy to certain extent but if combined with other tools as tumor markers and two dimensional ultrasounds, the sensitivity for malignancy increases.¹⁸ CT can be used to assess the severity of the disease in female with ovarian disorders. There is no strong evidence that CT is more specific and sensitive to find out ovarian cancer and USG is enough to evaluate the simple ovarian cysts. Jeong et al.³ showed that morphological characteristics associated with strong probability of malignancy were the presence of solid component (63%), papillary projection (92%), and free fluid in peritoneal cavity (56%).²⁰ Onyeka et al. found the sensitivity of CT scan for all ovarian cancer detection greater than that of USG 83% vs. 67%, but USG was more specific.²¹ In our study Overall, CT was found to have 98% sensitivity, 91% specificity, and an accuracy of 96% in the differentiation of benign and malignant ovarian masses, while PPV and NPV were 97% and 93%, respectively. The sensitivity of USG was 88%, specificity was 86% and PPV and NPV were 87% and 83% respectively. The findings of this study are corresponding to the results of Ahmed A et al.²² who found Trans Abdominal Sonography (TAS) to be 78% sensitive and 88.8% specific and CT to be 91% sensitive and 81.4% specific in evaluating benignity and malignancy in adnexal masses. While we are discordant with the results of USG in the study of Behtash N et al.²³ showing a sensitivity of 91.2% and specificity of 68.3%; there is close similarity in CT results of current study with them, showing 85.3% sensitivity and 56.1% specificity. Verit FF et al.²⁴ while evaluating the diagnostic accuracy of different techniques in diagnosis of ovarian tumours in premenopausal women, found USG to be 83% sensitive and 92% specific and CT to be 91% sensitive and 96% specific.

CONCLUSION

In this present study showed significant differences in the two methods i.e USG and CT. CT is showing more advantages regarding tumor localization, characterization. Hence CT can be advised if the unusual abnormalities were observed in routine USG scan in the diagnosis of ovarian masses.

REFERENCE

1. Karthikeyan B, Girija B, Pyadala N. Role of USG and CT in patients with ovarian masses. *International Journal of Contemporary Medicine Surgery and Radiology*. 2019;4(3):C193-C195.
2. Froyman W, Landolfo C, De Cock B, Wynants L, Sladkevicius P, Testa AC, et al.. Risk of complications in patients with conservatively managed ovarian tumours (IOTA5): a 2-year interim analysis of a multicentre, prospective, cohort study. *Lancet Oncol* 2019; 20: 448–58. doi: 10.1016/S1470-2045(18)30837-4.
3. Jeong YY, Outwater EK, Kang HK. Imaging evaluation of ovarian masses. *Radiographics*. 2000;20(1):1445–1470.
4. Aziz Z, Sana S, Saeed S, Akram M. Institution based tumor registry from Punjab: five year data based analysis. *J Pak Med Assoc*. 2003;53 (2):350–353.
5. Moro F, Zannoni GF, Arciuolo D, Pasciuto T, Amoroso S, Mascilini F, et al.. Imaging in gynecological disease (11): clinical and ultrasound features of mucinous ovarian tumors. *Ultrasound ObstetGynecol* 2017; 50: 261–70.
6. Pérez-López FR, Chedraui P, Troyano-Luque JM. Perianal post-menopausal incidental adnexal masses and the risk of sporadic ovarian malignancy: new insights and clinical management. *Gynecol Endocrinol*. 2010;26 (6):631–643.
7. Woodward PJ, Hosseinzadeh K, Saenger JS. Radiologic staging of ovarian carcinoma with pathologic correlation. *Radiographics*. 2004;24 (4):225–246.
8. Nam E, Kim Y, Kim J, et al. Kim. Diagnosis and staging of ovarian cancer: comparative values of PET/CT, Doppler US, CT, and MRI correlated with histopathologic analysis. *J Clin Oncol*. 2008;26(15S):5567.
9. Kinkel K, Lu Y, Mehdizade A, Pelte MF, Hricak H. Indeterminate ovarian mass at ultrasound: incremental value of second imaging test for characterization-meta analysis and Bayesian analysis. *Radiology*. 2005;236 (2):85–94.
10. Tsili AC, Tsampoulas C, Charisiadi A, et al. Adnexal masses: accuracy of detection and differentiation with multidetector computed tomography. *Gynecol Oncol*. 2008;110 (4):22–31.
11. Ovarian cysts. Office on Women's Health. November 19, 2014. Archived from the original on 29 June 2015. Retrieved, 2015
12. Joshi M, Ganesan K, Munshi HN, Ganesan S, Lawande A. Ultrasound of adnexal masses. In *Seminars in Ultrasound, CT and MRI*. WB Saunders. 2008;29(2):72-97.
13. Funt SA, Hann LE. Detection and characterization of adnexal masses. *Radiologic Clinics of North America*. 2002;40(3):591-608.
14. Devine C, Szklaruk J, Tamm EP. Magnetic resonance imaging in the characterization of pelvic masses. In

- Seminars in Ultrasound, CT and MRI. WB Saunders. 2005;26(3):172-204.
15. Bennett GL, Slywotzky CM, Giovanniello G. Gynecologic causes of acute pelvic pain: spectrum of CT findings. *Radiographics*. 2002;22(4):785-801.
 16. Mimoun C , Fritel X , Fauconnier A , Deffieux X , Dumont A , Huchon C. Epidemiology of presumed benign ovarian tumors. *Journal de gynécologie, obstétrique et biologie de la reproduction*, 2013; 42(8):722 -9.
 17. Helm William. Ovarian Cysts. Archived from the original on 7 September 2013. Retrieved, 2013.
 18. Levine D, Brown DL, Andreotti RF, Benacerraf B, Benson CB, Brewster WR, et al. Management of asymptomatic ovarian and other adnexal cysts imaged at US: Society of Radiologists in Ultrasound Consensus Conference Statement. *Radiology*. 2010; 256(3):943 -54.
 19. HE, Nguyen HN. The role of prophylactic oophorectomy in cancer prevention. *Gynecologic Oncology*. 2001 -2010; 55(3):38 -41. 7.
 20. Liu J, Xu Y, Wang J. Ultrasonography, computed tomography and magnetic resonance imaging for diagnosis of ovarian carcinoma. *Eur. J. Radiol*. 2007; 62(2):328 -334.
 21. Onyeka BA, Attalla A, Deemer H . Comparative diagnostic values of greyn scale USS versus CT scan in primary management of gynaecological pelvic mass with emphasis on ovarian detection and staging. *J. Obstet. Gynaecol*. 2001; 21(6):516 -9
 22. Ahmed A, Zamir S, Saghir NJ. Characterization of adnexal masses on trans abdominal ultrasonography and CT scan. *Ann Pak Inst Med Sci*. 2013;9(1):48–51.
 23. Behtash N, Rahmani M, Ghotbizadeh F, Karimi M, Zarchi AM. Ultrasonography and computed tomography for management of adnexal masses in Iranian patients with suspected ovarian cancer: results of a prospective study. *Asian Pacific Journal of Cancer Prevention*. 2009;10:201-04.
 24. Verit FF, Pehlivan M. Transvaginal ultrasound and computed tomography combined with ca-125 determinations in preoperative evaluation of ovarian masses in premenopausal women. *Harran Üniversitesi Tıp Fakültesi Dergisi*. 2007;4(2): 50–54.