

ORIGINAL RESEARCH

Thyroid dysfunction in patients with abnormal uterine bleeding

¹Dr. Swan Lata Das, ²Dr. Shalini Kumari, ³Dr. Abha Rani Sinha

^{1,2}Assistant Professor, ³Professor & HOD, Department of Obstetrics and Gynaecology, Sri Krishna Medical College and Hospital, Muzaffarpur, Bihar, India

Corresponding author

Dr. Shalini Kumari

Assistant Professor, Department of Obstetrics and Gynaecology, Sri Krishna Medical College and Hospital, Muzaffarpur, Bihar, India

Received: 15 January, 2023

Accepted: 22 February, 2023

ABSTRACT

Background: Thyroid disorders are more common in women than in men and in older adults compared with younger age groups. The present study was conducted to assess thyroid dysfunction in patients with abnormal uterine bleeding. **Materials & Methods:** 90 patients with abnormal uterine bleeding were enrolled. Assessment of serum triiodothyronine (T3), thyroxine (T4), and thyroid stimulating hormone (TSH) level was done. **Results:** Age group 18-27 years had 25, 28-37 years had 30 and 38-47 years had 35 patients. The difference was non-significant ($P > 0.05$). The level of triiodothyronine was normal in 80, increased in 6 and decreased in 4, level of thyroxine was normal in 82, increased in 5 and decreased in 3 and level of TSH was normal in 76, increased in 8 and decreased in 6 patients. The difference was significant ($P < 0.05$). Out of 90 patients, 76 were euthyroid, 8 had hypothyroidism and 6 had hyperthyroidism. Under hypothyroidism, 3 had menorrhagia, 2 had oligomenorrhoea, 1 had menometorrhagia, 1 had polymenorrhoea and 1 had hypomenorrhoea. Under hyperthyroidism, 2 had menorrhagia, 2 had oligomenorrhoea, 1 had menometorrhagia and 1 had polymenorrhoea. The difference was significant ($P < 0.05$). **Conclusion:** Abnormal uterine bleeding patients had menorrhagia, oligomenorrhoea, menometorrhagia and polymenorrhoea complaints.

Key words: Abnormal uterine bleeding.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

INTRODUCTION

Abnormal uterine bleeding (AUB) is one of the most common clinical presentations. It occurs in 10-20% of women between 15-50 years of age.¹ This may significantly affect the quality of life, results in time off work, lead to surgical interventions including hysterectomy, and ultimately have a significant impact on the health care system. The causes of abnormal uterine bleeding are polyp, adenomyosis, leiomyoma, malignancy and hyperplasia, coagulopathy, ovulatory disorders, endometrial causes, iatrogenic, not classified.²

Thyroid disorders are more common in women than in men and in older adults compared with younger age groups. Hypothyroidism is associated with a wide spectrum of reproductive disorders ranging from abnormal sexual development, menstrual irregularities, and infertility.³ The impact of hypothyroidism on the menstrual cycle has been identified since the 1950s and leads to changes in cycle length and blood flow. Subclinical hypothyroidism has been associated with occult menorrhagia before becoming symptomatic. The

prevalence of subclinical hypothyroidism is as high as 9.5% in women.⁴

Hyperthyroidism occurring before puberty has been reported to delay the onset of menses. In women of fertile age group, oligomenorrhoea and amenorrhoea are the commonest abnormalities associated with hyperthyroidism. These irregularities sometimes precede thyroid dysfunction.⁵ In the present times, subclinical hyper- and hypothyroidism can be diagnosed very early, whereas these would have passed unnoticed a few decades ago. Timely detection of thyroid disorder in patients presenting with menstrual disorders and their management can prevent surgical intervention like curettage and hysterectomy.⁶ The present study was conducted to assess thyroid dysfunction in patients with abnormal uterine bleeding.

MATERIALS & METHODS

The present study comprised of 90 women patients with abnormal uterine bleeding. All gave their written consent to participate in the study.

Demographic data such as name, age, etc. was recorded. A thorough general physical examination,

systemic examination, gynaecological, and pelvic examination was carried out. Parameters such as parity, menstrual history, onset and duration of complaints and amount of blood flow was recorded. Laboratory investigations such as assessment of hemoglobin, platelets, triiodothyronine (T3),

thyroxine (T4), and thyroid stimulating hormone (TSH) level were measured in all patients. All patients were subjected to ultrasound of abdomen and pelvis. Results were tabulated and subjected to statistical analysis. P value less than 0.05 was considered significant.

RESULTS

Table I: Distribution of patients based on age group

Age group (Years)	Number	P value
18-27	25	0.72
28-37	30	
38-47	35	

Table I shows that age group 18-27 years had 25, 28-37 years had 30 and 38-47 years had 35 patients. The difference was non-significant (P> 0.05).

Table II: Assessment of thyroid profile

Thyroid function	Normal	Increased	Decreased	P value
Tri-iodothyronine	80	6	4	0.02
Thyroxine	82	5	3	0.01
TSH	76	8	6	0.05

Table II, graph I shows that the level of tri-iodothyronine was normal in 80, increased in 6 and decreased in 4, level of thyroxine was normal in 82,

increased in 5 and decreased in 3 and level of TSH was normal in 76, increased in 8 and decreased in 6 patients. The difference was significant (P< 0.05).

Graph I: Assessment of thyroid profile




Table III: Patterns of abnormal uterine bleeding

Bleeding pattern	Hypothyroidism (8)	Hyperthyroidism (6)	P value
Menorrhagia	3	2	0.05
Oligomenorrhoea	2	2	
Menometorrhagia	1	1	
Polymenorrhoea	1	1	
Hypomenorrhoea	1	0	

Table III shows that out of 90 patients, 76 were euthyroid, 8 had hypothyroidism and 6 had

hyperthyroidism. Under hypothyroidism, 3 had menorrhagia, 2 had oligomenorrhoea, 1 had

menometorrhagia, 1 had polymenorrhoea and 1 had hypomenorrhoea. Under hyperthyroidism, 2 had menorrhagia, 2 had oligomenorrhoea, 1 had menometorrhagia and 1 had polymenorrhoea. The difference was significant ($P < 0.05$).

DISCUSSION

Menstrual disorders pose a huge burden on gynecology OPD, accounting for approximately 20% of attendance.⁷ Thyroid hormones play an important role in normal reproductive physiology through direct effects on the ovaries and indirectly by interacting with sex hormone-binding globulin. Thyroid dysfunction can lead to menstrual irregularities and infertility. In India, thyroid disorders are among the most common endocrine diseases.⁸

Onset of thyroid disorders increases with age, and it is estimated that 26 % of premenopausal and menopausal women are diagnosed with thyroid disease.⁹ Thyroid dysfunction may have profound effects on the female reproductive system. Both hypothyroidism and hyperthyroidism are associated with a variety of changes in reproductive function, including delayed onset of puberty, anovulatory cycles and abnormally high fetal wastage.^{10,11} The present study was conducted to assess thyroid dysfunction in patients with abnormal uterine bleeding.

We found that age group 18-27 years had 25, 28-37 years had 30 and 38-47 years had 35 patients. Thakur et al¹² in 79 patients with abnormal uterine bleeding, 67 (84.8%) were euthyroid, 11 (13.9%) were hypothyroid, and 1 (1.2%) was hyperthyroidism. The most common type of abnormal uterine bleeding was menorrhagia 34 (43%), followed by polymenorrhoea 23 (29%), oligomenorrhoea 13 (16.5%), menometorrhagia 6 (7.6%), metrorrhagia 2 (2.5%), and hypomenorrhoea 1 (1.3%). The maximum number of patients was between 20-25 years with the mean age of 31 years. Among hypothyroid, 7 (8.8%) had subclinical hypothyroidism and 4 (5%) had frank hypothyroidism.

We observed that the level of tri-iodothyronine was normal in 80, increased in 6 and decreased in 4, level of thyroxine was normal in 82, increased in 5 and decreased in 3 and level of TSH was normal in 76, increased in 8 and decreased in 6 patients. Subedi et al¹³ determined incidence of thyroid disorders in dysfunctional uterine bleeding and its correlation with menstrual patterns. The incidence of gynecological OPD attendance due to abnormal uterine bleeding was 3% and the incidence of thyroid dysfunction was 10.6% with hypothyroidism being the commonest (9.3%). The commonest menstrual pattern found was menorrhagia/polymenorrhoea in 8 percent. We observed that out of 90 patients, 76 were euthyroid, 8 had hypothyroidism and 6 had hyperthyroidism. Under hypothyroidism, 3 had menorrhagia, 2 had oligomenorrhoea, 1 had menometorrhagia, 1 had polymenorrhoea and 1 had

hypomenorrhoea. Under hyperthyroidism, 2 had menorrhagia, 2 had oligomenorrhoea, 1 had menometorrhagia and 1 had polymenorrhoea. Joshi et al¹⁴ found the prevalence of thyroid dysfunction in diagnosed cases of abnormal uterine bleeding. Out of the total cases of abnormal uterine bleeding, 15 (15.79%) had thyroid dysfunction. Among total cases, 80 (84.21%) were euthyroid. The mean age of the patients was 33 ± 8 years. Among thyroid dysfunction, 9 (60.0%) were hypothyroid, 4 (26.66 %) were subclinical hypothyroid, and 2 (13.33 %) were hyperthyroid.

Ajmani et al¹⁵ comprised of 100 women with menstrual complaints. The control group consisted of 50 women. Thyroid function tests, anti-TPO antibody estimation, and endometrial sampling were done in all patients. In patients with menstrual disorders, 44% had thyroid disorders in which subclinical hypothyroidism was prevalent in 20%, overt hypothyroidism in 14%, and overt hyperthyroidism in 8% of the women. Autoimmune thyroid antibodies were present in 30% patients of women with menstrual disorders. On endometrial sampling, hypothyroid patients mainly had proliferative endometrium (42.85%) whereas hyperthyroid had atrophic endometrium (60%).

CONCLUSION

Authors found that abnormal uterine bleeding patients had menorrhagia, oligomenorrhoea, menometorrhagia and polymenorrhoea complaints.

REFERENCES

1. Poppe K, Glinoe D. Thyroid autoimmunity and hypothyroidism before and during pregnancy. *Human Reprod Update*. 2003;9:149-161.
2. Koutras DA. Disturbances of menstruation in thyroid disease. *Ann NY Acad Sci*. 1997;816:280-284.
3. Sharma N, Sharma A. Thyroid profile in menstrual disorders. *JK Science*. 2012;14(1):14-17.
4. Albers JR, Hull SK, Wesley RM. Abnormal uterine bleeding. *Am Fam Phys*. 2004;69(8):1915-1934.
5. Bals-Pratsch M, Geyter D, Muller C, et al. Episodic variations of prolactin, thyroid-stimulating hormone, luteinizing hormone, melatonin and cortisol in infertile women with subclinical hypothyroidism. *Human Reprod*. 1997;12:896-904.
6. Barnard K, Frayne SM, Skinner KM, Sullivan LM. Health status among women with menstrual symptoms. *J Womens Health (Larchmt)*. 2003 Nov;12(9):911-9.
7. Doifode CD, Fernandes K. Study of thyroid dysfunction in patients with dysfunctional uterine bleeding. *Journal of Obstetric and Gynaecological Societies of India*. 2001;51(2):93-5.
8. Singh L, Agarwal CG, Chowdhary SR, Mehra P, Khare R. Thyroid profile in infertile women. *Journal of Obstetric and Gynaecological Societies of India*. 1990;40:248.
9. Kumar AHS, Saravanan S. A study of prevalence of thyroid disorders in patients with abnormal uterine bleeding. *Int J Reprod Contracept Obstet Gynecol*. 2017 Mar;6(3):1036-9.

10. Kattel P. Thyroid function test in abnormal uterine bleeding. *Nepal Journal of Obstetric and Gynaecology*. 2017 Jul-Dec;12(2):74-8.
11. Pilli GS, Sethi B, Dhadad AV, Mathur PR. Dysfunctional uterine bleeding. *Journal of Obstetric and Gynaecological Societies of India*. 2001;52(3):87-89.
12. Thakur M, Maharjan M, Tuladhar H, Dwa Y, Bhandari S, Maskey S, Bajracharya M. Thyroid Dysfunction in Patients with Abnormal Uterine Bleeding in a Tertiary Care Hospital: A Descriptive Cross-sectional Study. *JNMA: Journal of the Nepal Medical Association*. 2020 May;58(225):333.
13. Subedi S, Banerjee B, Manisha C. Thyroid disorders in women with dysfunctional uterine bleeding. *Journal of Pathology of Nepal*. 2016 Sep 24;6(12):1018-20.
14. Joshi BR, Rizal S, Subedi S. Thyroid Dysfunction in Patient with Abnormal Uterine Bleeding in a Tertiary Hospital of Eastern Nepal: A Descriptive Cross-sectional Study. *JNMA: Journal of the Nepal Medical Association*. 2021 Jul;59(239):635.
15. Ajmani NS, Sarbhai V, Yadav N, Paul M, Ahmad A, Ajmani AK. Role of thyroid dysfunction in patients with menstrual disorders in tertiary care center of walled city of Delhi. *The Journal of Obstetrics and Gynecology of India*. 2016 Apr;66(2):115-9.